

This label has been updated according to the re-evaluation decision for Abamectin, RVD2025-04. While users are encouraged to follow this updated label immediately, the previously approved label is valid until 2027-07-31 in accordance with the phase out period set out in the re-evaluation decision for Abamectin, RVD2025-04. This previously approved label will be provided upon request by emailing hc.pmra.info-arla.sc@canada.ca. In your email please include the product name and Registration number of the label you are requesting.

2025-4932
2025-12-05

| | | |
|-------|----------|-------------|
| GROUP | 6 | INSECTICIDE |
|-------|----------|-------------|

AVID® 1.9% EC MITICIDE/INSECTICIDE

MITICIDE/ INSECTICIDE

COMMERCIAL

EMULSIFIABLE CONCENTRATE

For Control of Labelled Pests in Greenhouse Ornamental, Greenhouse Vegetables (Peppers, Cucumbers, and Tomatoes), Outdoor Ornamental, and Boxwood.

ACTIVE INGREDIENT:

Abamectin..... 19 g/L

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN AND PETS**

WARNING



POISON

POTENTIAL SKIN SENSITIZER

REGISTRATION NO.: **24485**
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **1 Litre**

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario N1G 4Z3
Telephone: 1-877-964-3682

LABEL

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

3.0 TOXICOLOGICAL INFORMATION

Recommendations for Medical Treatment for Abamectin Acute Toxicity: Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness) and muscle tremors. Toxicity following accidental ingestion of the concentrate can be minimized by inducing vomiting under the direction of a physician within one half hour of exposure. If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels) as indicated by clinical signs, symptoms and measurements. In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND PETS.

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Potential skin sensitizer. Prolonged or frequently repeated exposure may cause allergic skin reactions in some

2025-4932
2025-12-05

individuals. Harmful if inhaled or absorbed through the skin. Do not breathe spray mist. Avoid contact with skin. May be fatal if swallowed.

RESTRICTED ENTRY INTERVAL (REI): DO NOT enter or allow entry into treated areas during the intervals specified in the following table:

| Crop/Use | Pre-Harvest Interval (PHI) | Postapplication Activity | REI |
|------------------------------|----------------------------|--------------------------|----------|
| Greenhouse cucumbers/peppers | 3 days | All activities | 12 hours |
| Greenhouse tomatoes | 1 day | All activities | 12 hours |
| Greenhouse ornamentals | N/A | All activities | 12 hours |

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

DO NOT apply using handheld mistblower/airblast or handheld fogging equipment.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear coveralls over long-sleeved shirt and long pants, chemical-resistant gloves, socks, chemical-resistant footwear and headgear during mixing, loading, application, clean-up and repair.

Mixer/loaders and applicators must wear either a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH-approved canister approved for pesticides.

For handheld application, wear eye, head and respiratory protection when applying above waist height, including overhead.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Wash splashes from skin IMMEDIATELY with plenty of water. Remove PPE immediately after handling this product. Wash the outside of gloves with soap and water before removing. As soon as possible, wash thoroughly and change into clean clothing. After spraying, wash hands and shower thoroughly with soap and water.

Heavily contaminated or drenched clothing and other absorbent materials contaminated with this product should be discarded. Do not reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Wear freshly laundered clothes daily.

Do not wear contaminated shoes.

Users should wash hands and exposed skin before eating, drinking, chewing gum, using tobacco, applying cosmetics or using the toilet.

While using product, do not eat, drink or use tobacco (including smoking).

6.0 ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms. Observe spray buffer zones and vegetative filter strips specified under DIRECTIONS FOR USE.

Toxic to birds and small wild mammals.

Toxic to bees. Bees may be exposed through direct spray, spray drift, and residues on leaves, pollen and nectar in flowering crops and weeds. Minimize spray drift to reduce harmful effects on bees in habitats close to the application site. **DO NOT** apply during the crop blooming period where indicated in crop-specific directions. Avoid applications when bees are foraging in the treatment area in ground cover containing blooming weeds. When using managed bees for pollination services, **DO NOT** apply during the crop blooming period. To further minimize exposure to pollinators, refer to the complete guidance "Protecting Pollinators during Pesticide Spraying - Best Management Practices" on the Health Canada website (www.canada.ca/pollinators). Follow use-specific and crop-specific directions for application timing.

Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland. Care should be taken when using this product in an integrated pest management program where users are relying on the presence of beneficial arthropods. To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. To reduce risk to aquatic organisms from runoff, a vegetative filter strip of at least 10 metres wide between the application area edge and adjacent, downhill aquatic habitats must be observed, as specified under DIRECTIONS FOR USE.

Greenhouse use: Toxic to bees and other beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). May harm bees and other beneficial arthropods, including those used in greenhouse production. Avoid application when bees or other beneficial arthropods are in the treatment area.

Do not apply this product directly to aquatic habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, ditches and wetlands), estuaries or marine habitats. Do not contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment and disposal of wastes.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Syngenta Canada Inc.

7.0 STORAGE

Store in a tightly closed container in a cool, dry place. Do not use or store near heat or open flame. Store this product away from food or feed. Open dumping is prohibited. Do not reuse empty container.

8.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean up of spills.

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

2025-4932
2025-12-05

| | | |
|-------|----------|-------------|
| GROUP | 6 | INSECTICIDE |
|-------|----------|-------------|

AVID® 1.9% EC MITICIDE/INSECTICIDE

MITICIDE/ INSECTICIDE

COMMERCIAL

EMULSIFIABLE CONCENTRATE

For Control of Labelled Pests in Greenhouse Ornamental, Greenhouse Vegetables (Peppers, Cucumbers, and Tomatoes), Outdoor Ornamental, and Boxwood.

ACTIVE INGREDIENT:

Abamectin..... 19 g/L

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN AND PETS**

WARNING



POISON

POTENTIAL SKIN SENSITIZER

REGISTRATION NO.: **24485**
PEST CONTROL PRODUCTS ACT

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario
N1G 4Z3
Telephone: 1-877-964-3682

PAMPHLET

Table of Contents

| | <u>Section Number</u> |
|---|-----------------------|
| NOTICE TO USER | 1.0 |
| FIRST AID | 2.0 |
| TOXICOLOGICAL INFORMATION | 3.0 |
| PRECAUTIONS..... | 4.0 |
| PERSONAL PROTECTIVE EQUIPMENT (PPE) | 5.0 |
| ENVIRONMENTAL PRECAUTIONS | 6.0 |
| STORAGE | 7.0 |
| DISPOSAL | 8.0 |
| PRODUCT INFORMATION | 9.0 |
| DIRECTIONS FOR USE | 10.0 |
| General Information | 10.1 |
| Ground Application | 10.2 |
| Spray Buffer Zones | 10.3 |
| CROP USE DIRECTIONS | 11.0 |
| Greenhouse Ornamentals | 11.1 |
| MINOR USES | 12.0 |
| Greenhouse Vegetables..... | 12.1 |
| Outdoor Ornamentals | 12.2 |
| RESISTANCE MANAGEMENT RECOMMENDATIONS | 13.0 |

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

3.0 TOXICOLOGICAL INFORMATION

Recommendations for Medical Treatment for Abamectin Acute Toxicity: Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness) and muscle tremors. Toxicity following accidental ingestion of the concentrate can be minimized by inducing vomiting under the direction of a physician within one half hour of exposure. If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels) as indicated by clinical signs, symptoms and measurements. In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND PETS.

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Potential skin sensitizer. Prolonged or frequently repeated exposure may cause allergic skin reactions in some

2025-4932
2025-12-05

individuals. Harmful if inhaled or absorbed through the skin. Do not breathe spray mist. Avoid contact with skin. May be fatal if swallowed.

RESTRICTED ENTRY INTERVAL (REI): DO NOT enter or allow entry into treated areas during the intervals specified in the following table:

| Crop/Use | Pre-Harvest Interval (PHI) | Postapplication Activity | REI |
|------------------------------|----------------------------|--------------------------|----------|
| Greenhouse cucumbers/peppers | 3 days | All activities | 12 hours |
| Greenhouse tomatoes | 1 day | All activities | 12 hours |
| Greenhouse ornamentals | N/A | All activities | 12 hours |

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

DO NOT apply using handheld mistblower/airblast or handheld fogging equipment.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear coveralls over long-sleeved shirt and long pants, chemical-resistant gloves, socks, chemical-resistant footwear and headgear during mixing, loading, application, clean-up and repair.

Mixer/loaders and applicators must wear either a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH-approved canister approved for pesticides.

For handheld application, wear eye, head and respiratory protection when applying above waist height, including overhead.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Wash splashes from skin IMMEDIATELY with plenty of water. Remove PPE immediately after handling this product. Wash the outside of gloves with soap and water before removing. As soon as possible, wash thoroughly and change into clean clothing. After spraying, wash hands and shower thoroughly with soap and water.

Heavily contaminated or drenched clothing and other absorbent materials contaminated with this product should be discarded. Do not reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Wear freshly laundered clothes daily.

Do not wear contaminated shoes.

Users should wash hands and exposed skin before eating, drinking, chewing gum, using tobacco, applying cosmetics or using the toilet.

While using product, do not eat, drink or use tobacco (including smoking).

6.0 ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms. Observe spray buffer zones and vegetative filter strips specified under DIRECTIONS FOR USE.

Toxic to birds and small wild mammals.

Toxic to bees. Bees may be exposed through direct spray, spray drift, and residues on leaves, pollen and nectar in flowering crops and weeds. Minimize spray drift to reduce harmful effects on bees in habitats close to the application site. **DO NOT** apply during the crop blooming period where indicated in crop-specific directions. Avoid applications when bees are foraging in the treatment area in ground cover containing blooming weeds. When using managed bees for pollination services, **DO NOT** apply during the crop blooming period. To further minimize exposure to pollinators, refer to the complete guidance "Protecting Pollinators during Pesticide Spraying - Best Management Practices" on the Health Canada website (www.canada.ca/pollinators). Follow use-specific and crop-specific directions for application timing.

Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland. Care should be taken when using this product in an integrated pest management program where users are relying on the presence of beneficial arthropods. To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. To reduce risk to aquatic organisms from runoff, a vegetative filter strip of at least 10 metres wide between the application area edge and adjacent, downhill aquatic habitats must be observed, as specified under DIRECTIONS FOR USE.

Greenhouse use: Toxic to bees and other beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). May harm bees and other beneficial arthropods, including those used in greenhouse production. Avoid application when bees or other beneficial arthropods are in the treatment area.

Do not apply this product directly to aquatic habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, ditches and wetlands), estuaries or marine habitats. Do not contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment and disposal of wastes.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Syngenta Canada Inc.

7.0 STORAGE

Store in a tightly closed container in a cool, dry place. Do not use or store near heat or open flame. Store this product away from food or feed. Open dumping is prohibited. Do not reuse empty container.

8.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean up of spills.

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

9.0 PRODUCT INFORMATION

AVID 1.9% EC Miticide/Insecticide is an emulsifiable concentrate containing 19 grams of abamectin per litre which, when mixed with water according to the Directions for Use, will control leafminers and spider mites on greenhouse ornamentals, greenhouse peppers, cucumbers and tomatoes.

10.0 DIRECTIONS FOR USE

10.1 General Information

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

DO NOT APPLY BY AIR.

A Vegetative Filter Strip (VFS) of at least 10 meters wide must be constructed and maintained.

The VFS is required between the application area edge and adjacent, downhill aquatic habitats to reduce risk to aquatic organisms from runoff. Aquatic habitats include, but are not limited to, lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, and estuaries. The VFS is to be composed of grasses and may also include shrubs, trees, or other vegetation. Both VFS and spray buffer zones, if applicable, must be observed. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

For greenhouse use: DO NOT allow releases, effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Syngenta Canada Inc. at 1-87-SYNGENTA (1-877-964-3682) for information before applying any tank mix that is not specifically recommended on this label.

Syngenta Canada Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) of all potential tank-mixes under all environmental conditions or for all crop varieties. Tank mixes that are not specifically listed on this label should be tested on a small area first, under local conditions and using standard practices, to confirm the tank mix is suitable for widespread application.

To determine the physical compatibility of this product with other products, use a jar test.

Always read and follow label directions including WALES mixing order.

10.2 Ground Application

Field sprayer application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. **DO NOT** apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) medium classification. Boom height must be 60 cm or less above the crop or ground.

Airblast application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. **DO NOT** direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. **DO NOT** apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

10.3 Spray Buffer Zones

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes,

streams, reservoirs and wetlands) and estuarine/marine habitats.

| Method of application | Crop | | Spray Buffer Zones (metres) Required for the Protection of: | | | | |
|-----------------------|---|--------------------|---|------------------|-------------------------------------|------------------|---------------------|
| | | | Freshwater Habitat of Depths: | | Estuarine/Marine Habitat of Depths: | | Terrestrial Habitat |
| | | | Less than 1 m | Greater than 1 m | Less than 1 m | Greater than 1 m | |
| Field sprayer | Outdoor ornamentals (non-specific), Boxwood | | 35 | 15 | 5 | 2 | 0 |
| Airblast | Outdoor ornamentals | Early growth stage | 60 | 50 | 35 | 25 | 0 |
| | | Late growth stage | 50 | 40 | 25 | 20 | 0 |

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for the application of this product with conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website.

2025-4932
2025-12-05

11.0 CROP USE DIRECTIONS

11.1 Greenhouse Ornamentals

| CROP | GREENHOUSE ORNAMENTALS |
|---------------------------------|---|
| PEST | Spider mites |
| RATE | 30 mL product/100 L water Apply, for example, in 2000 - 4000 L of water/ha. In volumes of water below 2000 L, use a minimum of 600 mL AVID 1.9% EC Miticide/Insecticide/ha. If more than 4000 L of water/ha are required for good plant coverage, apply a maximum rate of 1200 mL AVID 1.9% EC Miticide/Insecticide/ha. |
| APPLICATION METHOD | Ground application Mix with sufficient water and apply as a foliar spray to obtain uniform coverage |
| APPLICATION TIMING | Apply when mites first appear and repeat as necessary to maintain control. |
| RESTRICTED ENTRY INTERVAL (REI) | 12 hours |
| NOTES | Use in sufficient water volume to obtain uniform coverage. Do not apply more than 1200 mL or less than 600 mL AVID 1.9% EC Miticide/Insecticide per hectare. DO NOT apply this product through any type of irrigation system. Pollinator Protection: Toxic to bees and other beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). May harm bees and other beneficial arthropods, including those used in greenhouse production. Avoid application when bees or other beneficial arthropods are in the treatment area. |

2025-4932
2025-12-05

Page 16 of 20

| | |
|--|---|
| PEST | Leafminers |
| RATE | 60 mL product/100 L water Apply, for example, in 1000 - 2000 L of water/ha. In water volumes below 1000 L, use a minimum of 600 mL AVID 1.9% EC Miticide/Insecticide/ha. If more than 2000 L of water/ha are required for good plant coverage, apply the maximum rate of 1200 mL AVID 1.9%EC Miticide/Insecticide/ha. |
| APPLICATION METHOD | Ground application Mix with sufficient water and apply as a foliar spray to obtain uniform coverage |
| APPLICATION TIMING | Apply as necessary to maintain control but not at less than 7-day intervals. |
| RESTRICTED ENTRY INTERVAL (REI) | 12 hours |
| NOTES | Use in sufficient water volume to obtain uniform coverage. Do not apply more than 1200 mL or less than 600 mL AVID 1.9% EC Miticide/Insecticide per hectare. DO NOT apply this product through any type of irrigation system. Pollinator Protection: Toxic to bees and other beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). May harm bees and other beneficial arthropods, including those used in greenhouse production. Avoid application when bees or other beneficial arthropods are in the treatment area. |

12.0 MINOR USES

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Syngenta Canada Inc. under the User Requested Minor Use Label Expansion program. For these uses, Syngenta Canada Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

12.1 Greenhouse Vegetables

Mix with sufficient water and apply as a foliar spray to obtain uniform coverage. Apply when listed pests first appear and repeat as necessary to maintain control.

| GREENHOUSE VEGETABLES | |
|--|--|
| CROP | GREENHOUSE PEPPERS |
| PEST | Twospotted spider mite, Tomato psyllid (<i>Bactericerca [Paratrioza] cockerelli</i>) |
| RATE | 30 mL product/100 L water |
| APPLICATION METHOD | Ground application Mix with sufficient water and apply as a foliar spray to obtain uniform coverage |
| APPLICATION TIMING | Apply when pests first appear and repeat as necessary to maintain control. |
| RESTRICTED ENTRY INTERVAL (REI) | 12 hours |
| NOTES | Make no more than 5 applications per crop cycle. Apply no more than 6000 mL product/ha per crop cycle (114 g abamectin/ha per crop cycle). Application is limited to between February and October and/or when daily light intensity in the greenhouse is at levels higher than 700 Joules/cm ² /day. DO NOT apply within 3 days of harvest. |
| CROP | GREENHOUSE CUCUMBERS |
| PEST | Twospotted spider mite, Leafminer (<i>Liriomyza</i> spp.) |
| RATE | 30 mL product/ 100 L water |
| APPLICATION METHOD | Ground application Mix with sufficient water and apply as a foliar spray to obtain uniform coverage |
| APPLICATION TIMING | Apply when pests first appear and repeat as necessary to maintain control. |
| RESTRICTED ENTRY INTERVAL (REI) | 12 hours |
| NOTES | Make no more than 5 applications per crop cycle. Apply no more than 6000 mL product/ha per crop cycle (114 g abamectin/ha per crop cycle). DO NOT apply within 3 days of harvest. |

2025-4932
2025-12-05

Page 18 of 20

| | |
|--|--|
| CROP | GREENHOUSE TOMATOES |
| PEST | Twospotted spider mite, Leafminer (<i>Liriomyza</i> spp.), Tomato psyllid (<i>Bactericerca [Paratrioza] cockerelli</i>) |
| RATE | 30 mL product/ 100 L water |
| APPLICATION METHOD | Ground application Mix with sufficient water and apply as a foliar spray to obtain uniform coverage |
| APPLICATION TIMING | Apply when pests first appear and repeat as necessary to maintain control. |
| RESTRICTED ENTRY INTERVAL (REI) | 12 hours |
| NOTES | Apply no more than 3600 mL product/ha per crop cycle (68.4 g abamectin/ha per crop cycle). DO NOT apply within 1 day of harvest. |
| <p>Use in sufficient water volume to obtain uniform coverage. Application should be made preferably in 2000 - 4000 L of water/ha. If volumes of water below 2000 L must be used, apply a minimum of 600 mL AVID 1.9% EC Miticide/Insecticide /ha. If volumes of water greater than 4000 L of water/ha are required for good plant coverage, apply a maximum rate of 1200 mL AVID 1.9% EC Miticide/Insecticide /ha.</p> <p>Do not apply more than 1200 mL or less than 600 mL AVID 1.9% EC Miticide/Insecticide per hectare.</p> <p>DO NOT apply this product through any type of irrigation system.</p> <p>AVID 1.9% EC Miticide/Insecticide has been evaluated for phytotoxicity on a wide range of ornamental plants and on greenhouse peppers, cucumbers and tomatoes. However, since all combinations for sequences of pesticide sprays including surfactants and adjuvants have not been tested, it is recommended that a small area be sprayed first to make certain that no phytotoxicity occurs. Phytotoxicity has been observed following the use of AVID 1.9% EC Miticide/Insecticide on certain species of ferns (e.g. <i>Adiantum</i> spp.) and Shasta daisies (<i>Leucanthemum</i> spp.). It is, therefore, recommended that AVID 1.9% EC Miticide/Insecticide not be used on ferns or Shasta daisies.</p> <p>Pollinator protection: Toxic to bees and other beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). May harm bees and other beneficial arthropods, including those used in greenhouse production. Avoid application when bees or other beneficial arthropods are in the treatment area.</p> | |

12.2 Outdoor Ornamentals

| | |
|---|---|
| CROP | OUTDOOR ORNAMENTALS |
| PEST | Leafminers ^a |
| RATE | 600 – 1200 mL per ha (11.4 – 22.6 g a.i./ha) Use the higher rate when pest populations are high and/or when foliage is dense. |
| APPLICATION METHOD | Ground application Apply as a foliar spray to obtain uniform coverage |
| APPLICATION TIMING | Apply first application when leafminers first appear but before the economic threshold is exceeded. Consult provincial guidelines and local extension experts for optimal timing in your area. |
| MAXIMUM NUMBER OF APPLICATIONS PER YEAR | Do not apply more than 3 times per year |
| MAXIMUM AMOUNT OF ACTIVE INGREDIENT /HA/YEAR | DO NOT exceed a maximum cumulative application rate of 38.2 g a.i./ha/per year. |
| APPLICATION INTERVAL | 7 days |
| RESTRICTED ENTRY INTERVAL (REI) | Do not re-enter treated areas until residues have dried. |
| NOTES | ^a Apply in sufficient water volume to obtain uniform coverage; 1000 - 2000 L of water/ha is recommended. DO NOT apply using aerial application equipment. Pollinator Protection: Toxic to bees. DO NOT apply during the crop blooming period (onset of flowering until after petal fall is complete). (Excludes ornamental grasses and coniferous evergreens: pine, fir, juniper, spruce, arborvitae, hemlock, cypress, yew). |
| CROP | BOXWOOD |
| PEST | Boxwood psyllid ^b |
| RATE | 600 – 1200 mL per ha (11.4 – 22.6 g a.i./ha) Use the high rate when boxwood psyllid populations are high and/or when the boxwood canopy is dense. |
| APPLICATION METHOD | Ground application (field sprayer) Apply as a foliar spray to obtain uniform coverage |
| APPLICATION TIMING | Apply first application when psyllids first appear. Consult provincial guidelines and local extension experts for optimal timing in your area. |
| MAXIMUM NUMBER OF APPLICATIONS PER YEAR | Do not apply more than 3 times per year |
| MAXIMUM AMOUNT OF ACTIVE INGREDIENT /HA/YEAR | DO NOT exceed a maximum cumulative application rate of 38.2 g a.i./ha/per year. |
| APPLICATION INTERVAL | 7 days |
| RESTRICTED ENTRY INTERVAL (REI) | Do not re-enter treated areas until residues have dried. |
| NOTES | ^b Apply in sufficient water volume to obtain uniform coverage. DO NOT apply using aerial application equipment. Pollinator Protection: Toxic to bees. DO NOT apply during the crop blooming period. |

13.0 RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, please note that AVID 1.9% EC Miticide/Insecticide contains a Group 6 miticide/insecticide. Any insect population may contain individuals naturally resistant to AVID 1.9% EC Miticide/Insecticide and other Group 6 miticides/insecticides. The resistant individuals may dominate the insect population if this group of miticides/insecticides is used repeatedly in the same site. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay miticide/insecticide resistance:

Where possible, rotate the use of AVID 1.9% EC Miticide/Insecticide or other Group 6 miticides/insecticides with different groups that control the same pests.

It is recommended that AVID 1.9% EC Miticide/Insecticide not be applied more than twice in sequence or more than 6 times per crop (annually) or per year for perennials.

Use tank mixtures with miticides/insecticides from a different group when such use is permitted.

Miticide/insecticide use should be based on an IPM program that includes scouting and record keeping, and considers cultural, biological and other chemical control practices.

Monitor treated pest populations for resistance development.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

For further information or to report suspected resistance, contact company representatives at 1-877-SYNGENTA (1-877-964-3682) or at www.syngenta.ca.