

2023-6947
2024-07-24

RootShield® WP - Biological Fungicide

For the Suppression of Root Rot caused by Pythium spp., Rhizoctonia spp. and Fusarium spp. in Greenhouse Crops, Outdoor Nursery Plants, Greenhouse Transplants, Greenhouse and Field Herbs, Field Spices, Ginseng, Hops, and Cannabis (marihuana) produced commercially indoors.

COMMERCIAL
WETTABLE POWDER
READ THE LABEL BEFORE USING

ACTIVE INGREDIENT: *Trichoderma harzianum* Rifai strain KRL-AG2 contains a minimum of 1.0×10^7 colony forming units per gram of dry weight

Warning, contains the allergens wheat and sulfites.

POTENTIAL SENSITIZER

LIVE ORGANISMS

REGISTRATION No. 29890 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 0.45 – 90.9 kg

BioWorks, Inc. 100 Rawson Road, Suite 205, Victor, NY 14564
Telephone: 1.585.924.4362

Lot No.:

Date of Manufacture: Use within 12 months of date of manufacture.

GENERAL INFORMATION: RootShield® WP - Biological Fungicide is a biological fungicide for the suppression of root rot in greenhouse crops, outdoor nursery plants, greenhouse vegetable transplants, ginseng, hops, and Cannabis (marihuana) produced commercially indoors. The active ingredient is a beneficial microbe *Trichoderma harzianum* strain T-22 (KRL-AG2). When applied to transplants, greenhouse planting mix or soil, RootShield® WP - Biological Fungicide actively grows onto plant roots as they develop and protects roots against *Pythium*, *Rhizoctonia* and *Fusarium*. RootShield® WP - Biological Fungicide can be used alone or in conjunction with certain chemical fungicides; contact manufacturer or distributor for compatibility Technical Bulletin.

For control of seed rot and damping off of internally infected cuttings, use chemical fungicides or other appropriate measures in addition to RootShield® WP - Biological Fungicide.

Note: RootShield® WP - Biological Fungicide contains live spores of a beneficial microbe. RootShield® WP - Biological Fungicide becomes active when soil temperatures are above 10 degrees C. Control is not effective while soils remain cold and is more effective in neutral or acidic soils than in alkaline soils.

DIRECTIONS FOR USE:

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

DO NOT allow releases, effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters.

DRENCH APPLICATION TO POTTING MIX OR SOIL

Greenhouse Peppers, Tomatoes, Cucumbers and Eggplant

Greenhouse Ornamental Crops

Outdoor Nursery Crops

Greenhouse Vegetable Transplants

Greenhouse and Field Herbs

Field Spices

Asian water spinach (tung choy, kankon, rau muong)

Hops

Ginseng

Cannabis (Marihuana) produced commercially indoors

POTTING MIX: Suspend RootShield® WP - Biological Fungicide in sufficient water (eg. 30-45 g/100L) to achieve uniform application and apply at the rate of 55-110 g per cubic metre (loose) of greenhouse potting mix, soil or planting beds. Within the stated rate range, use higher rates when conditions favour disease development or high disease pressure is anticipated. RootShield® WP - Biological Fungicide can be applied through low pressure watering nozzles such as fan nozzles or other watering systems. Agitate to maintain suspension.

Cannabis (marihuana) produced commercially indoors: RootShield WP is applied to potting mix or soil as a drench prior to planting. Within a crop cycle, one re-application can be made as a soil drench after 8 to 12 weeks, as determined by disease and crop monitoring to assess the level of disease pressure or crop damage from the target diseases. Number of applications per crop cycle: 1-2. Maximum number of applications per year: Six (6).

NURSERY DRENCH:

Suspend RootShield® WP - Biological Fungicide in sufficient water (eg. 30 – 45 g/100 L) to achieve uniform application and apply at the rate of 55 – 110 g per cubic metre (loose) of nursery potting mix, soil or planting beds. Within the stated rate range, use higher rates when conditions favour

Crops	Disease	Pathogens	Rate	Spray volume	Control/Suppression
Greenhouse Ornamental Crops	Root rot	<i>Pythium</i> spp. <i>Rhizoctonia</i> spp. <i>Fusarium</i> spp.	55-110 g/m ³	30-45 g/100 L	Suppression
Outdoor Nursery Crops					
Greenhouse Vegetable Transplants					
Bulb vegetables					
Leafy vegetables					
Brassica vegetables					
Fruiting vegetables					
Cucurbit vegetables					
Asian water spinach (tung choy, kankon, rau muong)					
Greenhouse Peppers, Tomatoes, Cucumbers and Eggplant					
Greenhouse and Field Herbs (CG 25 ¹)			3-4 kg/ha	1000 litres/ha	
Field Spices (CG 26 ²)			3-4 kg/ha	1000 litres/ha	
Cannabis (marihuana) produced commercially indoors					
Ginseng			3-4 kg/ha	1000 litres/ha	
Hops			3-4 kg/ha	1000 litres/ha	
Ornamental Bulbs	Root rot	<i>Pythium</i> spp. <i>Rhizoctonia</i> spp. <i>Fusarium</i> spp.	Dip bulbs in 60 g/L suspension		Suppression

disease development or high disease pressure is anticipated RootShield® WP - Biological Fungicide can be applied through low pressure watering nozzles such as fan nozzles or other watering systems. Agitate to maintain suspension.

For Asian water spinach (tung choy, kankon, rau muong), a maximum of 6 applications can be made per year following an 8- to 12-week reapplication interval.

GINSENG, HOPS: Suspend RootShield® WP - Biological Fungicide Wettable Powder in 1,000 litres of water and apply at the rate of 3-4 kg per hectare to the soil surface immediately around the ginseng plants. Within the stated range, use higher rates when conditions favour disease development or high disease pressure is anticipated. Sufficient irrigation or rainfall of at least 5 mm should occur within 24 hours after application to move the active ingredient into the root zone. Agitate to maintain suspension.

DIP APPLICATION

Bulbs: All Ornamental Crops

Dip bulbs in suspension of 60 g/L RootShield® WP - Biological Fungicide prior to planting.

It is not possible to test all ornamental varieties, and some may show delayed growth. RootShield® WP - Biological Fungicide should first be tested on small lot of plants, if the response of the variety is not known, before using it for commercial scale production.

Bulb vegetables: chive, Chinese chive, daylily, Elegans hosta, fritillaria, garlic (bulb, great-headed, serpent), onion (bulb, green), kurrat, leek, lady's leek, wild leek, lily, onion (bulb, Chinese, green, macrostem, pearl, potato, tree, Welsh), shallot

Leafy vegetables: amaranth, arugula, cardoon, celery, Chinese celery, celtuce, chervil, chrysanthemum (edible-leaved, garland), corn salad, cress (garden, upland), dandelion, dock, endive, fennel, lettuce (head, leaf), orach, parsley, purslane (garden, winter), radicchio, rhubarb, spinach, New Zealand spinach, vine spinach, swiss chard

Brassica vegetables: broccoli (Chinese, raab), Brussels sprouts, cabbage, Chinese cabbage, cauliflower, cavolo broccoli, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens

Fruiting vegetables: cocona, eggplant (including African, pea, scarlet), groundcherry, martynia, okra, pepino, pepper (bell, non-bell), sunberry, tomatillo, tamarillo, goji berry, tomato (including bush, currant)

Cucurbit vegetables: Chinese waxgourd, citron melon, cucumber, gherkin, gourd (edible), momordica, muskmelon (cantaloupe, casaba, crenshaw, golden pershaw, honeydew, honey balls, mango Persian, pineapple, Santa Claus, snake), pumpkin, squash (winter, summer), watermelon

¹Crop Group 25 (**Herbs**): Agrimony, fresh leaves; Agrimony, dried leaves; Amla, fresh leaves; Amla, dried leaves; Angelica, fresh leaves; Angelica, dried leaves; Angelica, dahurian, fresh leaves; Angelica, dahurian, dried leaves; Applemint, fresh leaves; Applemint, dried leaves; Avarum, fresh leaves; Avarum, dried leaves; Balloon pea, fresh leaves; Balloon pea, dried leaves; Balm, fresh leaves; Balm, dried leaves; Barrenwort, fresh leaves; Barrenwort, dried leaves; Basil, fresh leaves; Basil, dried leaves; Basil, American, fresh leaves; Basil, American, dried leaves; Basil, Greek, fresh leaves; Basil, Greek, dried leaves; Basil, holy, fresh leaves; Basil, holy, dried leaves; Basil, lemon, fresh leaves; Basil, lemon, dried leaves; Basil, Russian, fresh leaves; Basil, Russian, dried leaves; Bay, fresh leaves; Bay, dried leaves; Bearberry, fresh leaves; Bearberry, dried leaves; Bisongrass, fresh leaves; Bisongrass, dried leaves; Blue mallow, fresh leaves; Blue mallow, dried leaves; Boneset, fresh leaves; Boneset, dried leaves; Borage, fresh leaves; Borage, dried leaves; Borage, Indian, fresh leaves; Borage, Indian, dried leaves; Burnet, fresh leaves; Burnet, dried leaves; Burnet, garden, fresh leaves; Burnet, garden, dried leaves; Burnet, salad, fresh leaves; Burnet, salad, dried leaves; Butterbur, fresh leaves; Butterbur, dried leaves; Calamint, fresh leaves; Calamint, dried leaves; Calamint, large-flower, fresh leaves; Calamint, large-flower, dried leaves; Calamint, lesser, fresh leaves; Calamint, lesser, dried leaves; Calendula, fresh leaves; Calendula, dried leaves; Caltrop, fresh leaves; Caltrop, dried leaves; Camomile (Chamomile), fresh leaves; Camomile (Chamomile) dried leaves; Camomile (Chamomile), German, fresh leaves; Camomile (Chamomile), German, dried leaves; Camomile (Chamomile), Roman, fresh leaves; Camomile (Chamomile), Roman, dried leaves; Caraway, fresh leaves; Caraway, dried leaves; Cat's claw, fresh leaves; Cat's claw, dried leaves; Catnip, fresh leaves; Catnip, dried leaves; Catnip, Japanese, fresh leaves; Catnip, Japanese, dried leaves; Celandine, greater, fresh leaves; Celandine, greater, dried leaves; Celandine, lesser, fresh leaves; Celandine, lesser, dried leaves; Celery, dried leaves; Centaury, fresh leaves; Centaury, dried leaves; Chaste tree, fresh leaves; Chaste tree, dried leaves; Chaste tree, Chinese, fresh leaves; Chaste tree, Chinese, dried leaves; Chervil, dried leaves; Chinese blackberry, fresh leaves; Chinese blackberry, dried leaves; Chinese foxglove, fresh leaves; Chinese foxglove, dried leaves; Chive, dried leaves; Chive, Chinese, dried leaves; Cicely, sweet, fresh leaves; Cicely, sweet, dried leaves; Cilantro, dried leaves; Clary, fresh leaves; Clary, dried leaves; Coriander, Bolivian, fresh leaves; Coriander, Bolivian, dried leaves; Coriander, Vietnamese, fresh leaves; Coriander, Vietnamese, dried leaves; Costmary, fresh leaves; Costmary, dried leaves; Creat, fresh leaves; Creat, dried leaves; Culantro, fresh leaves; Culantro, dried leaves; Curry leaf, fresh leaves; Curry leaf, dried leaves; Curryplant, fresh leaves; Curryplant, dried leaves; Cut leaf, fresh leaves; Cut leaf, dried leaves; Damiana, fresh leaves; Damiana, dried leaves; Dillweed, dried leaves; Dokudami, fresh leaves; Dokudami, dried leaves; Echinacea, fresh leaves; Echinacea, dried leaves; Epazote, fresh leaves; Epazote, dried leaves; Eucommia, fresh leaves; Eucommia, dried leaves; Evening primrose, fresh leaves; Evening primrose, dried leaves; Eyebright, fresh leaves; Eyebright, dried leaves; Fennel, common, fresh leaves; Fennel, common, dried leaves; Fennel, Florence, dried leaves; Fennel, Spanish, fresh leaves; Fennel, Spanish, dried leaves; Fenugreek, fresh leaves; Fenugreek, dried leaves; Feverfew, fresh leaves; Feverfew, dried

leaves; Field pennycress, fresh leaves; Field pennycress, dried leaves; Flowers, edible, fresh leaves, multiple species; Flowers, edible, dried leaves, multiple species; Fumitory, fresh leaves; Fumitory, dried leaves; Galbanum, fresh leaves; Galbanum, dried leaves; Galega, fresh leaves; Galega, dried leaves; Gambir, fresh leaves; Gambir, dried leaves; Geranium, fresh leaves; Geranium, dried leaves; Geranium, lemon, fresh leaves; Geranium, lemon, dried leaves; Geranium, rose, fresh leaves; Geranium, rose, dried leaves; Germander, golden, fresh leaves; Germander, golden, dried leaves; Goldenrod, European, fresh leaves; Goldenrod, European, dried leaves; Goldenseal, fresh leaves; Goldenseal, dried leaves; Gotu kola, fresh leaves; Gotu kola, dried leaves; Greater periwinkle, fresh leaves; Greater periwinkle, dried leaves; Guayusa, fresh leaves; Guyana, dried leaves; Gumweed, fresh leaves; Gumweed, dried leaves; Gymnema, fresh leaves; Gymnema, dried leaves; Gypsywort, fresh leaves; Gypsywort, dried leaves; Hawthorn, fresh leaves; Hawthorn, dried leaves; Heal-all, fresh leaves; Heal-all, dried leaves; Hemp nettle, fresh leaves; Hemp nettle, dried leaves; Honewort, fresh leaves; Honewort, dried leaves; Honeybush, fresh leaves; Honeybush, dried leaves; Horehound, fresh leaves; Horehound, dried leaves; Horsemint, fresh leaves; Horsemint, dried leaves; Horsetail, fresh leaves; Horsetail, dried leaves; Hyssop, fresh leaves; Hyssop, dried leaves; Hyssop, anise, fresh leaves; Hyssop, anise, dried leaves; Indian tobacco, fresh leaves; Indian tobacco, dried leaves; Ironwort, fresh leaves; Ironwort, dried leaves; Ivy, fresh leaves; Ivy, dried leaves; Jamaica dogwood, fresh leaves; Jamaica dogwood, dried leaves; Jasmine, fresh leaves; Jasmine dried leaves; Labrador tea, fresh leaves; Labrador tea, dried leaves; Lavender, fresh leaves; Lavender, dried leaves; Lemon verbena, fresh leaves; Lemon verbena, dried leaves; Lemongrass, fresh leaves; Lemongrass, dried leaves; Lovage, fresh leaves; Lovage, dried leaves; Love-in-a-mist, fresh leaves; Love-in-a-mist, dried leaves; Mamaki, fresh leaves; Mamaki, dried leaves; Marigold, fresh leaves; Marigold, dried leaves; Marigold, African, fresh leaves; Marigold, African, dried leaves; Marigold, Aztec, fresh leaves; Marigold, Aztec, dried leaves; Marigold, French, fresh leaves; Marigold, French, dried leaves; Marigold, Irish lace, fresh leaves; Marigold, Irish lace, dried leaves; Marigold, licorice, fresh leaves; Marigold, licorice, dried leaves; Marigold, Mexican mint, fresh leaves; Marigold Mexican mint, dried leaves; Marigold, signet, fresh leaves; Marigold, signet, dried leaves; Marjoram, fresh leaves; Marjoram, dried leaves; Marjoram, pot, fresh leaves; Marjoram, pot, dried leaves; Marjoram, sweet, fresh leaves; Marjoram, sweet, dried leaves; Marshmallow, fresh leaves; Marshmallow, dried leaves; Meadowsweet, fresh leaves; Meadowsweet dried leaves; Mint, fresh leaves; Mint, dried leaves; Mint, corn, fresh leaves; Mint, corn, dried leaves; Mint, Korean, fresh leaves; Mint Korean, dried leaves; Monarda, fresh leaves; Monarda, dried leaves; Moringa, fresh leaves; Moringa, dried leaves; Motherwort, fresh leaves; Motherwort, dried leaves; Mountainmint, fresh leaves; Mountainmint, dried leaves; Mountainmint, clustered, fresh leaves; Mountainmint, clustered, dried leaves; Mountainmint, hoary, fresh leaves; Mountainmint, hoary, dried leaves; Mountainmint, Virginia, fresh leaves; Mountainmint, Virginia, dried leaves; Mountainmint, whorled, fresh leaves; Mountainmint, whorled, dried leaves; Mugwort, fresh leaves; Mugwort, dried leaves; Mulberry, white, fresh leaves; Mulberry, white, dried leaves; Mullein, fresh leaves; Mullein, dried leaves; Mustard, hedge, fresh leaves; Mustard, hedge, dried leaves; Nasturtium, fresh leaves; Nasturtium, dried leaves; Nasturtium, bush, fresh leaves; Nasturtium, bush, dried leaves; Nasturtium, garden, fresh leaves; Nasturtium, garden, dried leaves; Nettle, stinging, fresh leaves; Nettle, stinging, dried leaves; Oregano, fresh leaves; Oregano, dried leaves; Oregano, Mexican, fresh leaves; Oregano, Mexican, dried leaves; Oregano, Puerto Rico, fresh leaves; Oregano, Puerto Rico, dried leaves; Oswego tea, fresh leaves; Oswego teat, dried leaves; Pandan leaf, fresh leaves; Pandan leaf, dried leaves; Pansy, fresh leaves; Pansy, dried leaves; Paracress, fresh leaves; Paracress dried leaves; Parsley, dried leaves; Partridge berry, fresh leaves; Partridge berry, dried leaves; Patchouli, fresh leaves; Patchouli, dried leaves; Pennyroyal, fresh leaves; Pennyroyal, dried leaves; Pepper leaf, black, fresh leaves; Pepper leaf, black, dried leaves; Peppermint, fresh leaves; Peppermint, dried leaves; Perilla, fresh leaves; Perilla, dried leaves; Pill bearing spurge, fresh leaves; Pill bearing spurge, dried leaves; Pipsissewa, fresh leaves; Pipsissewa, dried leaves; Plantain, common, fresh leaves; Plantain, common, dried leaves; Rooibos, fresh leaves; Rooibos, dried leaves; Rose, fresh leaves; Rose, dried leaves; Rosemary, fresh leaves; Rosemary, dried leaves; Sage, fresh leaves; Sage, dried leaves; Sage, Greek, fresh leaves; Sage, Greek, dried leaves; Sage, Spanish, fresh leaves; Sage, Spanish, dried leaves; Sage, white, fresh leaves; Sage, white, dried leaves; Savory, summer, fresh leaves; Savory, summer, dried leaves; Savory, winter, fresh leaves; Savory, winter, dried leaves; Senna, fresh leaves; Senna,

dried leaves; Siberian fir, fresh leaves; Siberian fir, dried leaves; Skullcap, fresh leaves; Skullcap, dried leaves; Small flower willow head, fresh leaves; Small flower willow head, dried leaves; Sorrel, fresh leaves; Sorrel, dried leaves; Sorrel, French, fresh leaves; Sorrel, French, dried leaves; Sorrel, garden, fresh leaves; Sorrel, garden, dried leaves; Southernwood, fresh leaves; Southernwood, dried leaves; Spearmint, fresh leaves; Spearmint, dried leaves; Spearmint, Scotch, fresh leaves; Spearmint, Scotch, dried leaves; Spilanthes, fresh leaves; Spilanthes, dried leaves; Spotted beebalm, fresh leaves; Spotted beebalm, dried leaves; St. John's Wort, fresh leaves; St. John's Wort, dried leaves; Stevia, fresh leaves; Stevia, dried leaves; Stoneroot, fresh leaves; Stoneroot, dried leaves; Swamp leaf, fresh leaves; Swamp leaf, dried leaves; Tansy, fresh leaves; Tansy, dried leaves; Tarragon, fresh leaves; Tarragon, dried leaves; Thuja, fresh leaves; Thuja, dried leaves; Thyme, fresh leaves; Thyme, dried leaves; Thyme, creeping, fresh leaves; Thyme, creeping, dried leaves; Thyme, lemon, fresh leaves; Thyme, lemon, dried leaves; Thyme, mastic, fresh leaves; Thyme, mastic, dried leaves; Toon, Chinese, fresh leaves; Toon, Chinese, dried leaves; Toothed clubmoss, fresh leaves; Toothed clubmoss, dried leaves; Trailing arbutus, fresh leaves; Trailing arbutus, dried leaves; Vasaka, fresh leaves; Vasaka, dried leaves; Verbena, blue, fresh leaves; Verbena, blue, dried leaves; Veronica, fresh leaves; Veronica, dried leaves; Violet, fresh leaves; Violet, dried leaves; Watermint, fresh leaves; Watermint, dried leaves; Waterpepper, fresh leaves; Waterpepper, dried leaves; Wild bergamot, fresh leaves; Wild bergamot, dried leaves; Wintergreen, fresh leaves; Wintergreen, dried leaves; Wood betony, fresh leaves; Wood betony, dried leaves; Woodruff, fresh leaves; Woodruff, dried leaves; Wormwood, fresh leaves; Wormwood, dried leaves; Wormwood, Roman, fresh leaves; Wormwood, Roman, dried leaves; Yarrow, fresh leaves; Yarrow, dried leaves; Yellow gentian, fresh leaves; Yellow gentian, dried leaves; Yerba santa, fresh leaves; Yerba santa, dried leaves; Yomogi, fresh leaves; Yomogi, dried leaves; Cultivars, varieties, and hybrids of these commodities.

²Crop Group 26 (Spices): Ajowan, seed; Alder buckhorn; Ambrette, seed; Angelica, dahurian, seed; Angelica, seed; Anise, seed; Autumn crocus; Barberry, bark; Betel vine; Birch, bark; Bisnaga, seed; Black bread weed; Bloodroot; Boldo, leaf; Calamus root (wild harvested); Caper buds; Caper spurge, seed; Caraway, black; Caraway, fruit; Cascara sagrada; Celery, seed; Chaste tree, berry; Chervil, seed; Chinese hawthorn; Chinese wineberry, fruit; Comfrey; Coptis; Coriander, fruit; Coriander, seed; Crampbark; Culantro, seed; Culvers root; Cumin; Cumin, black; Dill, seed; Echinacea, seed; Epimedium; Eucalyptus; Eucommia, bark; European beech; Felty germander; Fennel flower, seed; Fennel, common, fruit; Fennel, common, seed; Fennel, Florence, fruit; Fennel, Florence, seed; Fenugreek, seed; Galbanum, resin; Haw, black; Honewort, seed; Imperatoria; Indian tobacco, seed; Iva; Juniper berry; Linden, leaf; Lovage, seed; Magnolia, bark; Mahaleb; Milk thistle; Mioga; Mistletoe; Mojave yucca; Mustard, black; Mustard, brown; Mustard, seed; Mustard, white; Nasturtium, bush, pods; Nasturtium, garden, pods; Nasturtium, pods; Nettle, stinging, seed; Osha; Pepperbush, berry; Pepperbush, leaf; Perilla, seed; Phellodendron; Poppy, seed; Rue; Saffron crocus; Sassafras, bark; Sassafras, leaf; Sesame, seed; Skunk cabbage, root; Slippery elm; Sumac, fragrant; Sumac, smooth, leaf; Tasmanian pepper, berry; Tasmanian pepper, leaf; White willow; Willow; Witch hazel; Yaw root; Yellow gentian, roots; cultivars, varieties, and hybrids of these commodities.

GENERAL APPLICATION INSTRUCTIONS

RootShield® WP - Biological Fungicide can be applied using hand-held backpack or ground spray equipment. AGITATION IS REQUIRED TO MAINTAIN SUSPENSION. Application equipment should be cleaned before use of this product, and prepared sprays should be used within 4 hours of preparation. To prevent nozzle tip clogging the manufacturer recommends using 50 – 100 mesh inlet screens in all spray equipment. Within the stated application rate ranges, use higher rates when conditions favour disease development or high disease pressure is anticipated.

TANK MIXING

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner

labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Bioworks Inc. at 1-585-924-4362 for information before applying any tank mix that is not specifically recommended on this label.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.

May cause sensitization. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist. Workers and handlers (includes mixer/loader and applicators) must wear a long sleeved shirt, long pants, shoes plus socks, waterproof gloves and eye goggles as well as a dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C) or a NIOSH approved respirator with any N-95, R-95, P-95 or HE filter for biological products when mixing/loading or applying the product and until mists have settled and during all clean-up/repair activities. Wash thoroughly with soap and water after handling product. Remove contaminated clothing and follow manufacturer's directions for cleaning/maintaining personnel protective equipment (PPE) before use. If no such instructions are available use clothing detergent and hot water for cleaning all washable PPE. Keep and wash PPE separately from other laundry. Wash hands after use.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Can be applied up to and including the day of harvest. The pre-harvest interval is 0 days.

ENVIRONMENTAL PRECAUTIONS:

As this pesticide is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

DO NOT apply by air.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

GENERAL: Seek medical attention immediately if irritation or signs of toxicity occur and persist or is severe. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE:

Store in a secure dry place at temperatures between 2°C and 5°C. Do not contaminate water, food, or feed during storage. Keep product in original container during storage and keep container lid tightly closed when not in use. This product should be used within 12 months of the date of manufacture when stored under refrigerated conditions. Store this product away from food or feed.

DISPOSAL/ DECONTAMINATION:

1. Thoroughly empty the contents of the container into the application device or medium.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial/territorial requirements.
4. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.