

2025-2423
2025-11-20

GROUP	32	INSECTICIDE
--------------	-----------	--------------------

{Optional Marketing Claims for Container and Booklet in Brackets}
[container label]

SPEAR T LIQUID CONCENTRATE
Biological Insecticide/Acaricide

SOLUTION

Commercial

For Control or Suppression of Twospotted Spider Mites, Thrips and Whiteflies on Greenhouse Vegetables , Greenhouse Berries, Ornamentals grown in greenhouses and high tunnels and Cannabis grown in greenhouses or other enclosed structures, Spotted-wing Drosophila on Fruit Trees and Berries grown outdoors or in high tunnels and Berries grown in greenhouses, and aphids on Berries grown in greenhouses

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: GS-omega/kappa-Htx-Hv1a2.00%(w/w)

Contains 1,2-benzisothiazolin-3-one at 0.09% as a preservative

CAUTION-SKIN IRRITANT

REGISTRATION NO.: ___34679___ PEST CONTROL PRODUCTS ACT

NET CONTENTS: 0.5 L – 1050 L, Bulk

Batch/Lot:

Vestaron Corporation
4717 Campus Drive, Suite 1200
Kalamazoo, MI 49008, USA
Phone: 1-919-694-1022

2025-2423
2025-11-20

PRECAUTIONS:
KEEP OUT OF REACH OF CHILDREN

May cause skin irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE): Workers must wear long-sleeved shirt, long pants, chemical-resistant gloves, and socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab.

During airblast applications, workers must wear chemical-resistant coveralls with a chemical-resistant hood over long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH-approved canister approved for pesticides.

USER SAFETY RECOMMENDATIONS

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Restricted Entry Interval (REI): DO NOT enter or allow worker entry into treated areas for 4 hours, or until sprays have dried, unless wearing the appropriate PPE specified for the method of application.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Apply only in a way that this product will not contact workers or other persons, either directly or through drift. Only workers wearing personal protective equipment may be in the area where applications are occurring.

DO NOT apply by air.

FIRST AID

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

2025-2423
2025-11-20

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

STORAGE:

Store this product away from food or feed. Store product in original container at room temperature for up to one year. Do not allow product to freeze. Finish all the product after opening the bottle. Do not store opened bottle for later use.

DISPOSAL:

Recyclable containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable containers:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

2025-2423
2025-11-20

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2025-2423
2025-11-20

[booklet]

GROUP	32	INSECTICIDE
--------------	-----------	--------------------

SPEAR T LIQUID CONCENTRATE
Biological Insecticide/Acaricide

SOLUTION

Commercial

For Control or Suppression of Twospotted Spider Mites, Thrips and Whiteflies on Greenhouse Vegetables, Greenhouse Berries, Ornamentals grown in greenhouses and high tunnels and Cannabis produced grown in greenhouses or other enclosed structures, Spotted-wing Drosophila on Fruit Trees and Berries grown outdoors or in high tunnels and Berries grown in greenhouses, and against aphids on berries grown in greenhouses

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: GS-omega/kappa-Hctx-Hv1a2.00% (w/w)

Contains 1,2-benzisothiazolin-3-one at 0.09% as a preservative

CAUTION-SKIN IRRITANT

REGISTRATION NO.: ____34679____PEST CONTROL PRODUCTS ACT

NET CONTENTS: 0.5 L – 1050 L, Bulk

Vestaron Corporation
4717 Campus Drive, Suite 1200
Kalamazoo, MI 49008, USA
Phone: 1-919-694-1022

2025-2423
2025-11-20

PRECAUTIONS:
KEEP OUT OF REACH OF CHILDREN

May cause skin irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE): Workers must wear long-sleeved shirt, long pants, chemical-resistant gloves, and socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab.

During airblast applications, workers must wear chemical-resistant coveralls with a chemical-resistant hood over long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH-approved canister approved for pesticides.

USER SAFETY RECOMMENDATIONS

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Restricted Entry Interval (REI): DO NOT enter or allow worker entry into treated areas for 4 hours, or until sprays have dried, unless wearing the appropriate PPE as specified by the method of application.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Apply only in a way that this product will not contact workers or other persons, either directly or through drift. Only workers wearing personal protective equipment may be in the area where applications are occurring.

DO NOT apply by air.

FIRST AID

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

2025-2423
2025-11-20

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

STORAGE:

Store this product away from food or feed. Store product in original container at room temperature for up to one year. Do not allow product to freeze. Finish all the product after opening the bottle. Do not store opened bottle for later use.

DISPOSAL:

Recyclable containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable containers:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

2025-2423
2025-11-20

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean-up of spills.

PRODUCT INFORMATION

SPEAR T LIQUID CONCENTRATE is a biological insecticide/acaricide containing the peptide active ingredient GS-omega/kappa-Hctx-Hv1a for suppression of twospotted spider mites and thrips and control of whiteflies on greenhouse vegetables, ornamentals grown in greenhouses or high tunnels, and berries grown in greenhouses, suppression of twospotted spider mite on cannabis grown in greenhouses or other enclosed structures, suppression of aphids on berries grown in greenhouses, and suppression of spotted-wing drosophila on fruit trees and berries grown outdoors or in high tunnels and berries grown in greenhouses.

The active ingredient in SPEAR T LIQUID CONCENTRATE largely acts through contact and functions primarily as a central nervous system inhibitor of target pests infesting labeled crops. SPEAR T LIQUID CONCENTRATE is mixed with water and applied as a foliar spray.

DIRECTIONS FOR USE

SPEAR T LIQUID CONCENTRATE is effective on immature and adult life stages of the listed insects and mites. Close scouting and early attention to infestations is highly recommended. Earlier or younger developmental stages are more susceptible. {SPEAR T LIQUID CONCENTRATE is not effective on the egg stage.}

Thorough coverage of infested plant parts is necessary; SPEAR T LIQUID CONCENTRATE does not have systemic activity.

SPEAR T LIQUID CONCENTRATE has been evaluated for phytotoxicity on a variety of crops under various normal growing conditions. However, testing all crop varieties in all mixtures and combinations is not feasible. Prior to treating entire crop, test a small portion of the crop for sensitivity.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

DO NOT allow releases, effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters.

DO NOT apply by air.

DO NOT apply this product through any type of irrigation system.

2025-2423
2025-11-20

Integrated Pest Management (IPM):

SPEAR T LIQUID CONCENTRATE is an important tool in sound insect management whenever insecticide use is necessary. Apply SPEAR T LIQUID CONCENTRATE alone or in rotation with chemical insecticides. This will result in reduced susceptibility to insect damage, overall reduced use of chemical insecticides, and delayed development of pesticide resistance. Consult with a pest control advisor, extension agent or the manufacturer before treating an entire crop where beneficial insects serve as part of an IPM program. Consult local agricultural authorities for specific IPM strategies developed for your crop(s) and location.

Preharvest interval:

SPEAR T LIQUID CONCENTRATE can be applied up to and including the day of harvest. PHI=0 days

Application Instructions

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

Field sprayer application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (S572.1 to 572.3) Fine classification. Boom height must be 60 cm or less above the crop or ground.

Airblast application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. DO NOT apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

Apply as a foliar spray using conventional ground spray equipment (high pressure or low-volume sprayers). For indoor applications, mix 2-3 litres of SPEAR T with water to a total volume of 100 litres to prepare a 2 - 3% product solution. Apply in enough volume to achieve full coverage of the target crop (300-900 litres of this spray solution are generally sufficient to treat 1 hectare). For orchard crops, apply in a minimum spray volume of 470 L/ha. For other outdoor crops, apply in a minimum spray volume of 187 L/ha. For outdoor applications against spotted-wing drosophila, apply 9.4-28.0 litres of SPEAR T per hectare.

Use a spreader/sticker or non-ionic surfactant at 0.125% v/v to enhance the adhesion of SPEAR T LIQUID CONCENTRATE to the crop. Repeat applications at 3- to 10-day intervals depending upon plant growth rate, pest activity and climatic factors. Under heavy pest pressure conditions, use the shorter spray interval, the higher rate, and/or increase spray volume to improve spray coverage. Do not make more than three consecutive applications of SPEAR T LIQUID CONCENTRATE and do not make more than six applications per year. Do not spray to run off.

2025-2423
2025-11-20

PESTS CONTROLLED OR SUPPRESSED BY SPEAR T LIQUID CONCENTRATE ON THE FOLLOWING CROPS:

Vegetables Grown in Greenhouses

Crops	Insect/Mite Pests
Cucumber	Suppression of Twospotted Spider Mite (<i>Tetranychus urticae</i>)
Eggplant	
Pepper	Suppression of Thrips, includeing Western Flower Thrips (<i>Frankliniella occidentalis</i>)
Tomato	
Vegetable Transplants (Cucumbers, Tomatoes, Peppers and Eggplant)	Control of Whiteflies, includeing Sweetpotato Whitefly (<i>Bemisia tabaci</i>) and Greenhouse Whitefly (<i>Trialeurodes vaporariorum</i>)

2025-2423
2025-11-20

Herbaceous Ornamentals Grown in Greenhouses or High Tunnels

Crop	Insect/Mite Pests
Container grown flowering and foliage annuals and perennials Container or in-ground or soil grown annuals and perennials for cut flowers	Suppression of Twospotted Spider Mite (<i>Tetranychus urticae</i>) Suppression of Thrips, including Western Flower Thrips (<i>Frankliniella occidentalis</i>) Control of Whiteflies, including Sweetpotato Whitefly (<i>Bemisia tabaci</i>) and Greenhouse Whitefly (<i>Trialeurodes vaporariorum</i>)

Blueberry, Cherry and Caneberries (Crop Subgroup 13-07A) Grown Outdoors or in High Tunnels

Crop	Insect Pest
Blueberry, Cherry, and Caneberries (Crop Subgroup 13-07A) including blackberry (including Andean blackberry, arctic blackberry, bingleberry, black satin berry, boysenberry, brombeere, California blackberry, Chesterberry, Cherokee blackberry, Cheyenne blackberry, common blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, evergreen blackberry, Himalayaberry, hullberry, lavacaberry, loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, mora, mures deronce, nectarberry, Northern dewberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, ravenberry, rossberry, Shawnee blackberry, Southern dewberry, tayberry, youngberry, zarzamora, and cultivars, varieties and/or hybrids of these), loganberry, black and red raspberry, wild raspberry, and cultivars, varieties and/or hybrids of these.)	Suppression of Spotted-wing drosophila (<i>Drosophila suzukii</i>)

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Vestaron Corporation under the User Requested Minor Use Label Expansion program. For these uses, Vestaron Corporation has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

Caneberries (Crop Subgroup 13-07A) Grown in Greenhouses

Crop	Insect/Mite Pests
Caneberries (Crop Subgroup 13-07A) including blackberry (including Andean blackberry, arctic blackberry, bingleberry, black satin berry, boysenberry, brombeere, California blackberry, Chesterberry, Cherokee blackberry, Cheyenne blackberry, common blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, evergreen blackberry, Himalayaberry, hullberry, lavacaberry, loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, mora, mures deronce, nectarberry, Northern dewberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, Southern dewberry, tayberry, youngberry, zarzamora, and cultivars, varieties and/or hybrids of these), loganberry, black and red raspberry, wild raspberry, and cultivars, varieties and/or hybrids of these.	Suppression of Thrips, including Western Flower Thrips (<i>Frankliniella occidentalis</i>) Suppression of Spotted-wing Drosophila (<i>Drosophila suzukii</i>) Suppression of Twospotted Spider Mite (<i>Tetranychus urticae</i>) Suppression of Aphids Control of Whiteflies, including Sweetpotato Whitefly (<i>Bemisia tabaci</i>) and Greenhouse Whitefly (<i>Trialeurodes vaporariorum</i>)

Low Growing Berries (Crop Subgroup 13-07G) Grown in Greenhouses

Crop	Insect/Mite Pests
Low growing berries (Crop Subgroup 13-07G) including	Suppression of Thrips, including Western Flower

bearberry, bilberry, lowbush blueberry, cloudberry, cranberry, lingonberry, muntries, partridgeberry, strawberry, and cultivars, varieties and/or hybrids of these.	Thrips (<i>Frankliniella occidentalis</i>) Suppression of Spotted-wing Drosophila (<i>Drosophila suzukii</i>) Suppression of Twospotted Spider Mite (<i>Tetranychus urticae</i>) Suppression of Aphids Control of Whiteflies, including Sweetpotato Whitefly (<i>Bemisia tabaci</i>) and Greenhouse Whitefly (<i>Trialeurodes vaporariorum</i>)	
---	---	--

Cannabis Grown in Greenhouses or Other Enclosed Structures

Crop	Mite Pest
Cannabis grown in greenhouses or other enclosed structures	Suppression of Twospotted Spider Mite (<i>Tetranychus urticae</i>)

Foliar applications must only be made during the vegetative growth stage.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Vestaron Corporation at 919-694-1022 for information before applying any tank mix that is not specifically recommended on this label.

RESISTANCE MANAGEMENT

For resistance management, SPEAR T LIQUID CONCENTRATE contains a Group 32 insecticide/acaricide. Any insect population may contain individuals naturally resistant to SPEAR T LIQUID CONCENTRATE and other Group 32 insecticides/acaricides. The resistant individuals may dominate the insect population if this group of insecticides is used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed. To delay insecticide resistance:

- Where possible, rotate the use of SPEAR T LIQUID CONCENTRATE or other Group 32 insecticides/acaricides with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that is effective on the target pest when such use is permitted.

2025-2423
2025-11-20

- Insecticide/acaricide use should be based on an IPM program that includes scouting and record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development. Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

For further information and to report suspected resistance, contact Vestaron Corporation company representatives at 919-694-1022.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

[Optional Marketing Claims on Container or in Booklet]

{Total greenhouse advantage}
{For use against} {Controls or Suppresses} {Thrips} {Whiteflies} {Twospotted Spider Mites} in Greenhouses
{Control or Suppression of the top three major greenhouse pests: thrips, whiteflies and Twospotted spider mites}
{Biological insecticide for greenhouse thrips/whiteflies/Twospotted spider mites}
{For greenhouse {thrips} {whiteflies} {and} {twospotted spider mites} management}
{Labeled for use on food and non-food crops, ornamental flowers}
{
{For greenhouse use}
{Not for residential use}
{Requires dilution with water prior to application - see booklet for instructions}
{Spear® is a registered trademark of Vestaron Corporation}
{Vestaron name and logo are registered trademarks of Vestaron Corporation}
{Biological Insecticide}
{Biological Pesticide}
{Biological Miticide}
{Biological Insecticide and Miticide}
{Biological Peptide Insecticide}
{Biological Peptide Pesticide}
{Biological Peptide Miticide}
{Novel} {Unique} class of pesticides}
{Known mode of action}
{Novel} {Unique} mode of action}
{Peptide-based pesticide}
{Peptide-based insecticide}
{Peptide-based miticide} {For use with ornamentals}
{Liquid concentrate - easy to mix}
{No noxious odour}

2025-2423
2025-11-20

- {Storable at room temperature}
- {Zero-day pre-harvest interval}
- {For pre-harvest use}
- {May be sprayed just prior to harvest}
- {May be applied to open blooms}
- {Allowed for use over open bloom}
- {Compatible with integrated pest management practices}
- {Manufactured by fermentation}
- {Contains no live organisms}
- {Contains no organisms}
- {